



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

metatarsal of *Picoides* is present, though concealed beneath the skin, as noted by Beddard.

The 'Osteology of the Striges' is not up to the standard of the paper on the Woodpeckers, for the descriptive matter is too long and too much scattered to be readily used; moreover we lack the summary of characters at the end, this not being compensated for by the fact that the main characters are given on two or three pages at the outset. — F. A. L.

Chapman on the Genus *Sturnella*.¹—With the aid of a much larger series of specimens than has been available to previous writers on the subject, Mr. Chapman has been able to furnish the most valuable contribution to the history of this perplexing genus that has yet appeared. After characterizing the various races of Meadowlarks hitherto described, and pointing out the striking differences between the arid (*neglecta*) and the humid (*magna*) groups, the author considers the question of inter-relationships between the two. His conclusions are that the two groups were originally evolved in Mexico, *neglecta* occupying the table land and *magna* the lower humid areas, their relationships here being simply that of well marked geographic races.

Subsequently the two spread northward, *neglecta* covering the western United States and *magna* the eastern region. Still later the two forms, which in the north were quite distinct, have converged and intermingled, thus accounting for the presence of typical *magna* and *neglecta*, as well as intermediates or 'hybrids', side by side in Minnesota and Texas, a fact which is well established.

Due credit is given throughout the paper to Mr. E. W. Nelson, whose careful collecting in Mexico has cleared up the obscurity which had previously prevailed as to the relationships of the two forms in the south.

In emphasizing the fact that the main difference between the *magna* and *neglecta* groups is one of depth of color Mr. Chapman points out that the form recently described as *S. magna hoopesi* should be associated with the latter in spite of the extent of yellow on the sides of the throat. Unless it is deemed advisable to consider the above groups as representing two distinct species, no change in nomenclature is necessary.

Several interesting half-tones from photographs of skins and feathers illustrate the paper, and show clearly the contrast between the light and dark birds. — W. S.

Eaton's 'Birds of Western New York.'²—Mr. Eaton's list relates to "the western portion of New York State, extending eastward through the 'Finger-Lake' region," and consisting of three east and west belts,

¹ A Study of the Genus *Sturnella*. By Frank M. Chapman, Bull. Am. Mus. Nat. Hist., Vol. XIII, 1900, pp. 297-320.

² Birds of Western New York. By Elon Howard Eaton. Proc. Rochester Acad., Vol. IV, pp. 1-64. Feb. 15, 1901.